

Hydrophobic, pellicular, **monolithic** capillary columns based on cross-linked polynorbornene for biopolymer separations.

Mayr B, Hölzl G, Eder K, Buchmeiser MR, Huber CG.

Anal Chem. 2002 Dec 1;74(23):6080-7.

PMID:12498205[PubMed - indexed for MEDLINE]

- Capillary array high-performance liquid chromatography of  
2. **nucleic** acids and proteins.

Premstaller A, Oefner PJ, Oberacher H, Huber CG.

Anal Chem. 2002 Sep 15;74(18):4688-93.

PMID:12349971[PubMed - indexed for MEDLINE]

- Analysis of polymerase chain reaction products by on-line liquid  
3. chromatography-mass spectrometry for genotyping of  
polymorphic short tandem repeat loci.

Oberacher H, Parson W, Muhlmann R, Huber CG.

Anal Chem. 2001 Nov 1;73(21):5109-15.

PMID:11721907[PubMed - indexed for MEDLINE]

- High-performance liquid chromatography-electrospray ionization  
4. mass spectrometry of single- and double-stranded **nucleic**  
acids using **monolithic** capillary columns.

Premstaller A, Oberacher H, Huber CG.

Anal Chem. 2000 Sep 15;72(18):4386-93.

PMID:11008774[PubMed - indexed for MEDLINE]

- Specific applications of capillary electrochromatography to  
5. biopolymers, including proteins, **nucleic** acids, peptide  
mapping, antibodies, and so forth.

Krull IS, Sebag A, Stevenson R.

J Chromatogr A. 2000 Jul 28;887(1-2):137-63. Review.

PMID:10961309[PubMed - indexed for MEDLINE]